

Internal and External Loads of Young Elite Soccer Players during Defensive Small-Sided Games

by

Alberto Rabano-Muñoz^{1,*}, Luis Suarez-Arrones^{2,3}, Bernardo Requena³,
Jose A. Asian-Clemente^{1,2,3}

The aim of this study was to evaluate the effects of different time periods on the internal and external loads of soccer players during small-sided games (SSGs). Seventeen young soccer players performed a SSG of 5 vs. 5 + 5 with 2 floaters, where two teams had possession of the ball, and one had to recover it. With established periods of 30 s (SSG₃₀), 1 min (SSG₁) or 2 min (SSG₂), teams participated in a defensive role for these periods of time. Total distance covered, moderate speed running distance, high speed running distance, sprint running distance, accelerations, decelerations and Player Load were monitored using global positioning systems devices. The maximal heart rate and modified training impulse were monitored using heart rate monitors. The rating of perceived exertion (RPE) was also measured. Data showed a small increase between SSG₃₀ and SSG₁ in Player Load (ES = -0.35; $p < 0.01$), and a small increase in high speed running (ES = -0.41; $p < 0.05$) and sprinting (ES = -0.47; $p < 0.01$) between SSG₃₀ and SSG₂. Also, SSG₁ showed a small increase in sprinting (ES = -0.57; $p < 0.01$) and accelerations (ES = -0.37; $p < 0.05$) with respect to SSG₂. In addition, SSG₂ showed a small increase in the RPE with respect to SSG₃₀ (ES = 0.46; $p < 0.05$). The results indicate that shorter defensive periods in SSGs increased high speed running, while longer defensive periods promoted greater perceived exertion. Manipulation of the duration of defensive periods in SSGs is a variable that should be considered in soccer training.

Keywords: small-sided games; global positioning system (GPS); external load; heart rate; internal load

Introduction

Small-sided games (SSGs) are among the tasks most often used by soccer coaches during training (Halouani et al., 2014; Sarmiento et al., 2018). SSGs represent modified games played on reduced pitch areas, with a smaller number of players than competition matches, and adapted rules (Owen et al., 2020). The popularity of these soccer drills has grown in recent years due to their suitability for developing technical, tactical and physical aspects in a soccer-specific context at the same time (Jones and Drust, 2007; Owen et al., 2014), increasing motivation, and improving efficiency during training sessions (Casamichana et al., 2012). Furthermore, SSGs induce an integrated approach for soccer training because

they present advantages in terms of specificity of movements and inclusion of decision making (Hammami et al., 2018), and they allow to increase variability between sessions, ensuring that game movement patterns are replicated, as well as the physiological and technical demands of competition (Clemente et al., 2021).

In order to reproduce competition demands, multiple formats of SSGs have been developed and reported in the scientific literature (Aguar et al., 2012; Hill-Haas et al., 2011; Sarmiento et al., 2018). Some of the most popular variables investigated are pitch size, the number of players and the multiple combinations of them (Casamichana and Castellano, 2010; Kelly and Drust, 2009), inclusion of goalkeepers (Mallo and Navarro, 2008),

¹ Performance Department, Real Betis Balompié, Seville, Spain.

² Department of Sport Sciences, Universidad Pablo de Olavide, Seville, Spain.

³ Performance Department, Football Science Institute, Granada, Spain.

* Correspondence: albrm7@gmail.com

inclusion of floaters, numerical imbalances (Lacome et al., 2018; Rabano-Munoz et al., 2019), and encouragement from coaches (Rampinini et al., 2007). Another key variable during the design of an SSG is the duration format, which includes the lengths of the SSG, the rest intervals between sets, and the number of sets. SSG duration is extensively manipulated by coaches in their daily training, and has been shown to be a determinant factor for players' loads (Halouani et al., 2014; Sarmiento et al., 2018). Several previous studies have shown that long and continuous SSG duration formats generated greater player internal loads (Clemente et al., 2019a; Hill-Haas et al., 2009; Tessitore et al., 2006) in contrast to intermittent and short SSG formats, which showed greater total distance covered and more high speed running (Casamichana et al., 2013; Clemente et al., 2019a; Fanchini et al., 2011). Clemente et al. (2019a) analysed the variations in internal and external loads in a 5 vs. 5 SSG format with two different duration routines (3 x 6 min and 6 x 3 min), suggesting that longer sets increase the perception of effort, and contribute to a large decrease in running distances. Casamichana et al. (2013) used the same format of 5 vs. 5 SSG in a continuous format of 16 min, a 4 x 4 min format and a 2 x 8 min format, concluding that short duration formats resulted in a greater total distance covered and higher stability in physiological responses.

However, more research is still needed with regard to duration formats in SSGs. Previous studies have been carried out with players participating in defensive and offensive phases of the game for variable and uncontrolled periods. Considering that the behaviour of players is different in the defensive and offensive phases (Silva et al., 2014), and tactical decisions modulate physical actions employed by players (Schuth et al., 2016), defensive roles during the game could be more demanding than offensive roles (Castellano et al., 2022). Thereby, it is very frequent in soccer to design defensive SSGs where players must be in a defending mode for a specified period of time. However, although they are extensively used, to the best of our knowledge the consequences of these defensive drills and the responses of soccer players to them are still unknown. For this reason, the aim of this study was to evaluate the effects of specific time periods of defensive phase play on the internal and external loads of soccer players during

the same SSG. The hypothesis was that the duration of the defensive period would affect the external and internal loads of players. Furthermore, shorter defensive periods would be more demanding in external loads and longer defensive periods would increase internal loads.

Methods

Participants

Seventeen young soccer players (age: 15.2 ± 0.3 years; body height: 171.4 ± 6.5 cm; body mass: 62.5 ± 7.5 kg; YYIRT1 performance: 2365 ± 467 m) from an elite academy of a Spanish first division club participated in this study. Players participated in four training sessions and one competitive match (Sunday) per week, and were highly familiarized with SSGs as these were very frequently used in their daily training. All players who participated in the study were outfield players. These data were created as a condition of monitoring in which players' activities were measured during the competitive season (Winter and Maughan, 2009), so that the ethics committee clearance was not required. The study conformed nevertheless to the recommendations of the Declaration of Helsinki. Participants were informed of the design and gave their consent before the start of the study.

Measures

External loads were monitored using global positioning system devices operating at a sampling frequency of 10 Hz (Catapult Vector S7, Catapult Sports, Australia). The reliability and accuracy of these devices have been validated previously (Johnston et al., 2014). Variables measured were: total distance covered, distance covered between 14.0 and 17.9 km·h⁻¹ (moderate speed running), distance covered between 18.0 and 20.9 km·h⁻¹ (high speed running), and distance covered above 21.0 km·h⁻¹ (sprinting). These thresholds have been widely used previously (Asian-Clemente et al., 2022; Casamichana et al., 2012; Suarez-Arrones et al., 2015). Simultaneously, players' accelerations and decelerations above 2.5 m·s⁻² were recorded. This threshold has been previously used for medium and high intensity accelerations (Bradley et al., 2010; Dalen et al., 2021). Player Load was also measured. This variable is based on the acceleration data that are recorded by triaxial accelerometers integrated in the GPS devices and

have been used to assess the neuromuscular load (Gomez-Carmona et al., 2019; Reche-Soto et al., 2019). Each player wore a heart rate monitor (Polar Team 2®, Polar Electro Oy, Finland) to obtain values for the maximal heart rate (HR_{max}) reached during the SSG and the internal training load using the modified training impulse ($TRIMP_{mod}$) proposed by Stagno et al. (2007). In order to determine the individual HR_{max} of each player, all the players completed a YYIRT1 before the study (Bangsbo et al., 2008). The rate of perceived exertion (RPE) was also measured (Foster et al., 2001). All players were fully familiarized with the use of the RPE before the beginning of the study.

Design and Procedures

Data were collected from six training sessions during the 2020–2021 season (October–November). Measurements took place on the hardest day of the microcycle, when a higher neuromuscular load on the field was planned (Wednesday). One SSG was performed each training session. These sessions were performed on an artificial turf at the same time each day (16:00–18:00 pm) and begun with a 20-min standardized warm-up of mobility, active stretching and the same format and duration of passing drills.

The SSG format used in the study was 5 vs. 5 + 5 with 2 floaters (Figure 1), where two teams had to maintain ball possession, and one team had to recover it. When the defensive team intercepted the ball, or if during possession the ball left the field of play, a new ball was introduced by the technical staff, and the defensive team continued in the defensive role for the remaining duration of the task. The aim of the defensive team was to recover the highest number of balls during their defensive participation, while the offensive teams had to maintain ball possession. Depending on the defined period of 30 s (SSG_{30}), 1 min (SSG_1) or 2 min (SSG_2), teams took a defensive role for these periods of time and changed roles when the defined period was completed. The SSG was played using a continuous bout of 12 min on a 30 x 30 m pitch. Each team completed one period of the defensive role and two periods of the offensive role continuously until finishing the bout. The area per player was of 75 m² to ensure defensive technical-tactical aspects (Clemente and Sarmento, 2020; Fradua et al., 2013). SSG_{30} was performed on weeks 1 and 4, SSG_1 on weeks 2 and 5, and SSG_2 on weeks 3 and 6.

Statistical Analysis

Data are presented as mean \pm standard deviation (SD). The Shapiro-Wilk test demonstrated that all variables presented a normal distribution. Differences between players were determined using one-way analysis of variance (ANOVA). Post-hoc tests were calculated using Bonferroni correction for multiple comparisons to determine significant differences. The level of statistical significance was set at $p \leq 0.05$. Standardized differences in effect size (ES, 95% confidence interval [95%CI]) in selected variables were calculated. The threshold values for assessing magnitudes of the ES (changes as a fraction or multiple of baseline standard deviation) were < 0.20 , 0.20, 0.60, 1.2 and 2.0 for trivial, small, moderate, large and very large ES, respectively (Hopkins et al., 2009). SPSS (version 19, SPSS Inc., Chicago, IL, USA) was used for all statistical analyses.

Results

Table 1 shows the internal and external loads for the studied SSGs. Data analysis is presented in Figure 2, and revealed a small increase between SSG_{30} and SSG_1 in Player Load (ES = -0.35 [-0.53 – -0.17]; $p < 0.01$). A small increase between SSG_{30} and SSG_2 was observed in high speed running (ES = -0.41 [-0.76 – -0.06]; $p < 0.05$) and sprinting (ES = -0.47 [-0.79 – -0.15]; $p < 0.01$). Between SSG_1 and SSG_2 , a small increase was shown in sprinting (ES = -0.57 [-0.90 – -0.24]; $p < 0.01$) and accelerations (ES = -0.37 [-0.67 – -0.07]; $p < 0.05$). In addition, SSG_2 displayed a small increase in the RPE compared to SSG_{30} (ES = 0.46 [0.12–0.80]; $p < 0.05$). Comparison of the rest of variables yielded unclear differences. Figure 3 shows the HR responses in the drills played.

Discussion

The aim of this study was to evaluate the effects of different fixed time periods during the defensive phase on the internal and external loads of soccer players, during the same SSG. The results are partially in line with the hypotheses. Even though players' responses during both SSG_{30} and SSG_1 were very similar, players showed a small increase in the RPE and a small decrease in high speed running distance and sprinting distance in SSG_2 .

It has been determined that the duration of SSGs can influence the external load of soccer players (Halouani et al., 2014; Sarmiento et al., 2018). Previous studies have shown that shorter duration induces greater total distance covered and more distance covered at high speed in SSGs (Casamichana et al., 2013; Clemente et al., 2019a; Hill-Haas et al., 2009). For example, Hill-Haas et al. (2009) compared a 1 x 24 min format with a 4 x 6 min format for three different SSGs, finding that players covered a significantly greater distance at high speed running in formats with shorter bouts. The present study found a small increase in high speed running distance during SSG₃₀ and SSG₁ in comparison with SSG₂. However, total distance did not change in the different formats analysed. Several processes could explain these differences in players' time-motion responses. The level of neuromuscular fatigue experienced by players during sets of 2 min may have been greater than during 1 min and 30 s bouts. It has been shown that fatigue during intense periods of match play is possibly related to intramuscular acidosis or the accumulation of potassium in the muscle interstitium (Mohr et al., 2005). It is possible that during SSG₂ players experienced similar processes to those experienced in some intense periods of match play. Also, the longer periods proposed for

SSG₃₀ and SSG₁ in which players continue playing in an offensive role (1 min and 2 min, respectively), but with more attackers (12 attackers vs. 5 defenders) could induce an active recovery and may be sufficient to facilitate recovery for involvement in high speed running actions. During matches, soccer players generally have a mean recovery time between high speed running bouts of 65–75 s (Bradley et al., 2009). This period of time has been shown to be sufficient for PCr re-synthesis and muscle re-oxygenation in repeated sprint studies (Brocherie et al., 2015; Padulo et al., 2015). On the other hand, a longer time in the defensive phase could induce a pacing strategy in players, resulting in less external loads due to awareness of the length of the defensive period and consequent self-organization of the pace (Sampson et al., 2015). By contrast, in shorter defensive play phases, the change of the role from offensive to defensive may induce aggressive behaviour related to ball recovery, and thus greater external loads. This behaviour may be strengthened by the numerical superiority of the offensive team, meaning that the defensive team may be working as a unit and waiting for the right moment to apply pressure (Nunes et al., 2020).

Table 1. Comparison of external and internal loads in the SSG studied.

	Mean ± SD			p value		
	SSG ₃₀	SSG ₁	SSG ₂	SSG ₃₀ vs. SSG ₁	SSG ₃₀ vs. SSG ₂	SSG ₁ vs. SSG ₂
DC (m)	1345.0 ± 115.4	1314.0 ± 148.9	1292.8 ± 159.4	0.402	0.099	0.596
MSR (m)	219.9 ± 80.8	207.8 ± 69.6	167.8 ± 54.9	0.982	0.100	0.038
HSR (m)	77.4 ± 44.2	52.3 ± 25.5	40.8 ± 24.5	0.242	0.049	0.187
SPR (m)	20.8 ± 22.1	17.5 ± 13.7	9.7 ± 12.7	0.514	0.021	0.001
Accelerations (#)	10.2 ± 3.5	11.0 ± 3.4	9.0 ± 3.1	0.375	0.387	0.043
Decelerations (#)	6.0 ± 2.7	6.0 ± 2.1	5.9 ± 2.8	0.773	0.845	0.592
Player Load (au)	159.3 ± 17.7	146.5 ± 14.2	153.9 ± 28.9	0.003	0.272	0.500
TRIMP _{mod} (au)	31.8 ± 7.3	28.4 ± 7.5	29.0 ± 7.9	0.091	0.212	0.817
HR _{max} (bpm)	191.4 ± 6.7	192.3 ± 8.7	193.4 ± 7.3	0.819	0.490	0.530
RPE (au)	7.6 ± 0.8	7.8 ± 0.5	8.0 ± 0.7	0.371	0.031	0.101

SD = standard deviation; *DC* = total distance covered; *MSR* = moderate speed running distance; *HSR* = high speed running distance; *SPR* = sprinting distance; *HR* = heart rate; *max* = maximal; *RPE* = rate of perceived exertion; *au* = arbitrary units; *bpm* = beats per minute.




Figure 1. Graphical representation of the SSG studied.




Figure 2. Comparison of external and internal loads in the SSG studied.

DC = total distance covered; MSR = moderate speed running distance; HSR = high speed running distance; SPR = sprinting distance; TRIMPmod = modified training impulse; HR = heart rate; max = maximal; RPE = rate of perceived exertion; ES = effect size.




Figure 3. Comparison of the HR in the three fixed time periods of the defensive phase in the SSG format.

SSG = small-sided game; HR = heart rate.

With regard to internal load responses, previous investigations have shown contradictory results. While some studies showed that longer formats produced higher heart rate and RPE values in comparison with shorter duration formats (Clemente et al., 2019a; Hill-Haas et al., 2009; Tessitore et al., 2006), another found that heart rate responses appeared to remain higher in a 4 x 4 min SSG format than in a 2 x 8 or a 1 x 16 min game,

using a 5 vs. 5 SSG format (Casamichana et al., 2013). Our results showed no differences in TRIMP_{mod} and HR_{max} between SSGs when the defensive phase was 30 s, 1 min or 2 min. Probably, the minority transition between phases of play with specific periods of defence induced specific pacing strategies in the teams (Nunes et al., 2020), provoking a decrease in exercise intensity during this phase. Although our results did not show

statistically significant differences in TRIMP_{mod} and HR_{max} between the SSG formats carried out, a small increase in the RPE was recorded by players during SSG₂ in comparison with SSG₃₀. These data are in line with the findings of previous studies, showing greater RPE values in longer duration formats of SSGs in comparison to shorter duration formats (Clemente et al., 2019b; Hill-Haas et al., 2009). Although the external load was greater in SSG₃₀, and there were no differences in the internal training load in the three specified defence periods, neuromuscular fatigue effects may have been the cause of the increase in perceived exertion. This may apply specifically to SSG₂, where players had to maintain their efforts to recover the ball for 2 minutes, producing a multidimensional fatigue effect because they had to perform defensive decision-making actions continuously (Smits et al., 2014).

To our knowledge, this is the first study to evaluate the effects of defensive SSGs in soccer players. Although this study provides new insights for designing specific training, some limitations must be acknowledged. First, although the differences found are statistically significant, the values shown by the ES are small. This could be due to the size of the sample, or the high variability of external and internal loads during SSGs (Clemente et al., 2022). Therefore, future studies should take these aspects into account. Only a format with offensive superiority (5 vs. 5 + 5 players with 2 floaters) was investigated, thus these data cannot be extrapolated to other formats. Previous research indicates that high-inferiority numeric situations during SSGs may induce minor running demands. This may be due to the fact that in situations with a high level of difficulty related

to recovering the ball, players in the defensive phase perform conservative behaviours (Torres-Ronda et al., 2015). Future studies could analyse the consequences of defensive SSGs with numerical equality or lower numerical imbalances, and their suitability to reproduce the demands of soccer matches. Despite the fact that external and internal loads can be affected by the phase of play (Castellano et al., 2022), this study did not differentiate between the offensive and defensive phases. Future studies could focus on this issue to understand how the tactical context affects the training load. In addition, in this study there was no comparison made between defensive and traditional SSGs (where change of the phase is accomplished without using established periods), thus studies that investigate this comparison are needed.

The present investigation provides useful information for coaches about the duration of SSGs that constitute one of the main constraints in the daily design of training. Our results indicate that shorter fixed periods in defensive SSGs increased high speed running, while longer fixed periods resulted in greater perceived exertion. Manipulation of the time duration of defensive SSGs is another variable that should be considered during soccer training when these tasks are used. These findings suggest that practitioners could use shorter fixed periods in defensive SSGs with the aim to increase the external load of players in the strongest session of the microcycle. Furthermore, longer periods in defensive SSGs could be used to increase the fatigue effect of players. Therefore, coaches should avoid longer defensive periods in the days preceding the match.

Author Contributions: Conceptualization: A.R.-M. and J.A.A.-C.; methodology: A.R.-M. and J.A.A.-C.; software: L.S.-A. and B.R.; validation: A.R.-M., L.S.-A., B.R. and J.A.A.-C.; formal analysis: B.R. and J.A.A.-C.; investigation: A.R.-M. and J.A.A.-C.; resources: L.S.-A. and B.R.; data curation: A.R.-M. and J.A.A.-C.; writing—original draft preparation: A.R.-M. and J.A.A.-C.; writing—review & editing: L.S.-A., B.R. and J.A.A.-C.; visualization: A.R.-M.; supervision: L.S.-A., B.R. and J.A.A.-C.; project administration: A.R.-M., L.S.-A., B.R. and J.A.A.-C. All authors have read and agreed to the published version of the manuscript.

Funding Information: No sources of funding were used to assist in the preparation of this work.

Institutional Review Board Statement: This study was conducted following the principles of the Declaration of Helsinki. These data were created as a condition of monitoring in which player's activities were measured during the competitive season, so that the ethics committee clearance was not required.

Informed Consent: Informed consent was obtained from all participants included in the study.

Conflicts of Interest: The authors have no potential conflicts of interest that are directly relevant to the content of this study.

Acknowledgements: The authors of the study would like to thank the players who participated and the coaches for their compliance.

References

- Aguiar, M., Botelho, G., Lago, C., Macas, V., & Sampaio, J. (2012). A review on the effects of soccer small-sided games. *Journal of Human Kinetics*, 33, 103–113. doi: 10.2478/v10078-012-0049-x
- Asian-Clemente, J., Rabano-Munoz, A., Requena, B., & Suarez-Arrones, L. (2022). High-speed training in a specific context in soccer: transition games. *International Journal of Sports Medicine*, 43(10), 881–888. doi: 10.1055/a-1794-9567
- Bangsbo, J., Iaia, F. M., & Krstrup, P. (2008). The Yo-Yo intermittent recovery test: a useful tool for evaluation of physical performance in intermittent sports. *Sports Medicine*, 38(1), 37–51. doi: 10.2165/00007256-200838010-00004
- Bradley, P. S., Di Mascio, M., Peart, D., Olsen, P., & Sheldon, B. (2010). High-intensity activity profiles of elite soccer players at different performance levels. *Journal of Strength and Conditioning Research*, 24(9), 2343–2351. doi: 10.1519/JSC.0b013e3181aeb1b3
- Bradley, P. S., Sheldon, W., Wooster, B., Olsen, P., Boanas, P., & Krstrup, P. (2009). High-intensity running in English FA Premier League soccer matches. *Journal of Sports Sciences*, 27(2), 159–168. doi: 10.1080/02640410802512775
- Brocherie, F., Millet, G. P., & Girard, O. (2015). Neuro-mechanical and metabolic adjustments to the repeated anaerobic sprint test in professional football players. *European Journal of Applied Physiology and Occupational Physiology*, 115(5), 891–903. doi: 10.1007/s00421-014-3070-z
- Casamichana, D., & Castellano, J. (2010). Time-motion, heart rate, perceptual and motor behaviour demands in small-sides soccer games: effects of pitch size. *Journal of Sports Sciences*, 28(14), 1615–1623. doi: 10.1080/02640414.2010.521168
- Casamichana, D., Castellano, J., & Castagna, C. (2012). Comparing the physical demands of friendly matches and small-sided games in semiprofessional soccer players. *Journal of Strength and Conditioning Research*, 26(3), 837–843. doi: 10.1519/JSC.0b013e31822a61cf
- Casamichana, D., Castellano, J., & Dellal, A. (2013). Influence of different training regimes on physical and physiological demands during small-sided soccer games: continuous vs. intermittent format. *Journal of Strength and Conditioning Research*, 27(3), 690–697. doi: 10.1519/JSC.0b013e31825d99dc
- Castellano, J., Errekagorri, I., Los Arcos, A., Casamichana, D., Martin-Garcia, A., Clemente, F. M., Lopez-Del Campo, R., Resta, R., & Echeazarra, I. (2022). Tell me how and where you play football and I'll tell you how much you have to run. *Biology of Sport*, 39(3), 607–614. doi: 10.5114/biolsport.2022.106155
- Clemente, F. M., Afonso, J., & Sarmiento, H. (2021). Small-sided games: An umbrella review of systematic reviews and meta-analyses. *PLoS One*, 16(2), e0247067. doi: 10.1371/journal.pone.0247067
- Clemente, F. M., Aquino, R., Praca, G. M., Rico-Gonzalez, M., Oliveira, R., Silva, A. F., Sarmiento, H., & Afonso, J. (2022). Variability of internal and external loads and technical/tactical outcomes during small-sided soccer games: a systematic review. *Biology of Sport*, 39(3), 647–672. doi: 10.5114/biolsport.2022.107016
- Clemente, F. M., Nikolaidis, P. T., Rosemann, T., & Knechtle, B. (2019a). Variations of Internal and External Load Variables between Intermittent Small-Sided Soccer Game Training Regimens. *International Journal of Environmental Research and Public Health*, 16(16), 2923. doi: 10.3390/ijerph16162923
- Clemente, F. M., & Sarmiento, H. (2020). The effects of small-sided soccer games on technical actions and skills: A systematic review. *Human Movement*, 21(3), 100–119. doi: 10.5114/hm.2020.93014
- Clemente, F. M., Sarmiento, H., Rabbani, A., Van Der Linden, C., Kargarfard, M., & Costa, I. T. (2019b). Variations of external load variables between medium- and large-sided soccer games in professional players. *Research in Sports Medicine*, 27(1), 50–59. doi: 10.1080/15438627.2018.1511560

- Dalen, T., Sandmael, S., Stevens, T. G. A., Hjelde, G. H., Kjosnes, T. N., & Wisloff, U. (2021). Differences in Acceleration and High-Intensity Activities Between Small-Sided Games and Peak Periods of Official Matches in Elite Soccer Players. *Journal of Strength and Conditioning Research*, 35(7), 2018–2024. doi: 10.1519/JSC.0000000000003081
- Fanchini, M., Azzalin, A., Castagna, C., Schena, F., McCall, A., & Impellizzeri, F. M. (2011). Effect of bout duration on exercise intensity and technical performance of small-sided games in soccer. *Journal of Strength and Conditioning Research*, 25(2), 453–458. doi: 10.1519/JSC.0b013e3181c1f8a2
- Foster, C., Florhaug, J. A., Franklin, J., Gottschall, L., Hrovatin, L. A., Parker, S., Doleshal, P., & Dodge, C. (2001). A new approach to monitoring exercise training. *Journal of Strength and Conditioning Research*, 15(1), 109–115.
- Fradua, L., Zubillaga, A., Caro, O., Ivan Fernandez-Garcia, A., Ruiz-Ruiz, C., & Tenga, A. (2013). Designing small-sided games for training tactical aspects in soccer: extrapolating pitch sizes from full-size professional matches. *Journal of Sports Sciences*, 31(6), 573–581. doi: 10.1080/02640414.2012.746722
- Gomez-Carmona, C. D., Pino-Ortega, J., Sanchez-Urena, B., Ibanez, S. J., & Rojas-Valverde, D. (2019). Accelerometry-Based External Load Indicators in Sport: Too Many Options, Same Practical Outcome? *International Journal of Environmental Research and Public Health*, 16(24), 5101. doi: 10.3390/ijerph16245101
- Halouani, J., Chtourou, H., Gabbett, T., Chaouachi, A., & Chamari, K. (2014). Small-sided games in team sports training: a brief review. *Journal of Strength and Conditioning Research*, 28(12), 3594–3618. doi: 10.1519/JSC.0000000000000564
- Hammami, A., Gabbett, T. J., Slimani, M., & Bouhlel, E. (2018). Does small-sided games training improve physical fitness and team-sport-specific skills? A systematic review and meta-analysis. *Journal of Sports Medicine and Physical Fitness*, 58(10), 1446–1455. doi: 10.23736/S0022-4707.17.07420-5
- Hill-Haas, S. V., Dawson, B., Impellizzeri, F. M., & Coutts, A. J. (2011). Physiology of small-sided games training in football: a systematic review. *Sports Medicine*, 41(3), 199–220. doi: 10.2165/11539740-000000000-00000
- Hill-Haas, S. V., Rowsell, G. J., Dawson, B. T., & Coutts, A. J. (2009). Acute physiological responses and time-motion characteristics of two small-sided training regimes in youth soccer players. *Journal of Strength and Conditioning Research*, 23(1), 111–115. doi: 10.1519/jsc.0b013e3181818efc1a
- Hopkins, W. G., Marshall, S. W., Batterham, A. M., & Hanin, J. (2009). Progressive statistics for studies in sports medicine and exercise science. *Medicine and Science in Sports and Exercise*, 41(1), 3–13. doi: 10.1249/MSS.0b013e3181818cb278
- Johnston, R. J., Watsford, M. L., Kelly, S. J., Pine, M. J., & Spurr, R. W. (2014). Validity and interunit reliability of 10 Hz and 15 Hz GPS units for assessing athlete movement demands. *Journal of Strength and Conditioning Research*, 28(6), 1649–1655. doi: 10.1519/JSC.0000000000000323
- Jones, S., & Drust, B. (2007). Physiological and technical demands of 4 v 4 and 8 v 8 in elite youth soccer players. *Kinesiology*, 39(2), 150–156.
- Kelly, D. M., & Drust, B. (2009). The effect of pitch dimensions on heart rate responses and technical demands of small-sided soccer games in elite players. *Journal of Science and Medicine in Sport*, 12(4), 475–479. doi: 10.1016/j.jsams.2008.01.010
- Lacome, M., Simpson, B. M., Cholley, Y., & Buchheit, M. (2018). Locomotor and Heart Rate Responses of Floaters During Small-Sided Games in Elite Soccer Players: Effect of Pitch Size and Inclusion of Goalkeepers. *International Journal of Sports Physiology and Performance*, 13(5), 668–671. doi: 10.1123/ijsp.2017-0340
- Mallo, J., & Navarro, E. (2008). Physical load imposed on soccer players during small-sided training games. *Journal of Sports Medicine and Physical Fitness*, 48(2), 166–171.
- Mohr, M., Krstrup, P., & Bangsbo, J. (2005). Fatigue in soccer: a brief review. *Journal of Sports Sciences*, 23(6), 593–599. doi: 10.1080/02640410400021286
- Nunes, N. A., Goncalves, B., Coutinho, D., & Travassos, B. (2020). How Numerical Unbalance Constraints Physical and Tactical Individual Demands of Ball Possession Small-Sided Soccer Games. *Frontiers in Psychology*, 11, 1464. doi: 10.3389/fpsyg.2020.01464

- Owen, A. L., Newton, M., Shovlin, A., & Malone, S. (2020). The Use of Small-Sided Games as an Aerobic Fitness Assessment Supplement Within Elite Level Professional Soccer. *Journal of Human Kinetics*, 71, 243–253. doi: 10.2478/hukin-2019-0086
- Owen, A. L., Wong, D. P., Paul, D., & Dellal, A. (2014). Physical and technical comparisons between various-sided games within professional soccer. *International Journal of Sports Medicine*, 35(4), 286–292. doi: 10.1055/s-0033-1351333
- Padulo, J., Tabben, M., Ardigo, L. P., Ionel, M., Popa, C., Gevat, C., Zagatto, A. M., & Dello Iacono, A. (2015). Repeated sprint ability related to recovery time in young soccer players. *Research in Sports Medicine*, 23(4), 412–423. doi: 10.1080/15438627.2015.1076419
- Rabano-Munoz, A., Asian-Clemente, J., Saez de Villarreal, E., Nayler, J., & Requena, B. (2019). Age-Related Differences in the Physical and Physiological Demands during Small-Sided Games with Floaters. *Sports*, 7(4), 79. doi: 10.3390/sports7040079
- Rampinini, E., Impellizzeri, F. M., Castagna, C., Abt, G., Chamari, K., Sassi, A., & Marcora, S. M. (2007). Factors influencing physiological responses to small-sided soccer games. *Journal of Sports Sciences*, 25(6), 659–666. doi: 10.1080/02640410600811858
- Reche-Soto, P., Cardona-Nieto, D., Diaz-Suarez, A., Bastida-Castillo, A., Gomez-Carmona, C., Garcia-Rubio, J., & Pino-Ortega, J. (2019). Player Load and Metabolic Power Dynamics as Load Quantifiers in Soccer. *Journal of Human Kinetics*, 69, 259–269. doi: 10.2478/hukin-2018-0072
- Sampson, J. A., Fullagar, H. H., & Gabbett, T. (2015). Knowledge of bout duration influences pacing strategies during small-sided games. *Journal of Sports Sciences*, 33(1), 85–98. doi: 10.1080/02640414.2014.925571
- Sarmiento, H., Clemente, F. M., Harper, L. D., Costa, I. T., Owen, A., & Figueiredo, A. J. (2018). Small sided games in soccer—a systematic review. *International Journal of Performance Analysis in Sport*, 18(5), 693–749. doi: <https://doi.org/10.1080/24748668.2018.1517288>
- Schuth, G., Carr, G., Barnes, C., Carling, C., & Bradley, P. S. (2016). Positional interchanges influence the physical and technical match performance variables of elite soccer players. *Journal of Sports Sciences*, 34(6), 501–508. doi: 10.1080/02640414.2015.1127402
- Silva, B., Garganta, J., Santos, R., & Teoldo, I. (2014). Comparing Tactical Behaviour of Soccer Players in 3 vs. 3 and 6 vs. 6 Small-Sided Games. *Journal of Human Kinetics*, 41, 191–202. doi: 10.2478/hukin-2014-0047
- Smits, B. L., Pepping, G. J., & Hettinga, F. J. (2014). Pacing and decision making in sport and exercise: the roles of perception and action in the regulation of exercise intensity. *Sports Medicine*, 44(6), 763–775. doi: 10.1007/s40279-014-0163-0
- Stagno, K. M., Thatcher, R., & van Someren, K. A. (2007). A modified TRIMP to quantify the in-season training load of team sport players. *Journal of Sports Sciences*, 25(6), 629–634. doi: 10.1080/02640410600811817
- Suarez-Arrones, L., Torreno, N., Requena, B., Saez De Villarreal, E., Casamichana, D., Barbero-Alvarez, J. C., & Munguia-Izquierdo, D. (2015). Match-play activity profile in professional soccer players during official games and the relationship between external and internal load. *Journal of Sports Medicine and Physical Fitness*, 55(12), 1417–1422.
- Tessitore, A., Meeusen, R., Piacentini, M. F., Demarie, S., & Capranica, L. (2006). Physiological and technical aspects of "6-a-side" soccer drills. *Journal of Sports Medicine and Physical Fitness*, 46(1), 36–43.
- Torres-Ronda, L., Goncalves, B., Marcelino, R., Torrents, C., Vicente, E., & Sampaio, J. (2015). Heart Rate, Time-Motion, and Body Impacts When Changing the Number of Teammates and Opponents in Soccer Small-Sided Games. *Journal of Strength and Conditioning Research*, 29(10), 2723–2730. doi: 10.1519/JSC.0000000000000932
- Winter, E. M., & Maughan, R. J. (2009). Requirements for ethics approvals. *Journal of Sports Sciences*, 27(10), 985. doi: 10.1080/02640410903178344

Received: 04 October 2022; Accepted: 07 December 2022; Published: 20 April 2023